

- ◆ One adult should stay with the child/young person to assess the severity of symptoms and in case of:

- Itchiness
- Tingling of lips and face
- Tummy cramps
- Vomiting
- Blotchiness of skin

Give \_\_\_\_\_ (Oral Antihistamine) \_\_\_ml at once

In cases of:

- Wheeziness
- Swelling of face and throat
- Difficulty in breathing/swallowing
- Feeling faint

Place child/young person on floor in recovery position (Safe Airway Position)

Give preloaded adrenaline injection to outer thigh  
(this can be administered through light clothing).

- ◆ If no breathing/pulse, initiate basic life support (CPR).
- ◆ If there is no improvement to above action within 10 minutes and there are symptoms of weakness/floppiness pallor then:

Repeat preloaded adrenaline injection once more if 2<sup>nd</sup> preloaded adrenaline injection is available

- ◆ Hand over child/young person's care to Ambulance Team/parents on their arrival
- ◆ Handover preloaded adrenaline injection to ambulance staff or if this hasn't been done, safely dispose of it.
- ◆ Record all medication given with date and time of administration

#### **Awareness**

The headteacher will arrange for the staff in the setting to be briefed about his/her condition and about other arrangements contained in this document.

The setting staff will take all reasonable steps to ensure that \_\_\_\_\_ (Name) does not eat any food items unless they have been prepared/approved by his/her parents.

\_\_\_\_\_ (Name) parents will remind their child regularly of the need to refuse any food items, which might be offered to them by other children/young people.

In particular, \_\_\_\_\_ (Name) parents will provide for him/her the following food items: \_\_\_\_\_